

ALTA DIAGNOSTICS, INC. 3123 Research Way Ste 214, Carson City NV 89706 (800) 359-9691 (775) 283-5780 FAX: (775) 283-5787

18 MONTH

OPEN VIAL

STABILITY

LIQUID URINE CONTROL

FOR

MICROSCOPIC & HIGH SPECIFIC GRAVITY

| | LOT #041519 POSITIVE Exp 03/17 | | LOT #041520 NEGATIVE Exp 03/17 | | PROCEDURE |
|------------------|-----------------------------------|---------------------|-----------------------------------|---------------------|--|
| | MICROSCOPIC | SPECIFIC GRAVITY | MICROSCOPIC | SPECIFIC GRAVITY | |
| HIGH | 35 | 1.039 | | 1.039 | 1. Shake well before using to assure |
| MEAN | 20 CELL/HP ± 15 | 1.034 ± .005 | 0 CELL/HP | 1.034 ± .005 | complete mixing of the contents. |
| LOW | 5 | 1.029 | | 1.029 | 2. Remove bottle cap and pour 12 ml into a clean, dry conical centrifuge tube.* |
| DAY 1 | | | | | |
| DAY 2 | | | | | 3. Centrifuge for 5 minutes at 2000 rpm. (A lower rpm may be used if this is called for in your laboratory procedure. However, a somewhat lower mean may result!) |
| DAY 3 | | | | | |
| DAY 4 | | | | | |
| DAY 5 | | | | | 4. Remove control from the centrifuge and at this time, if desired, take and record the specific gravity reading by placing a small urinometer in the centrifuge tube or, alternatively, transfer a few drops of the supernate to a refractometer. |
| DAY 6 | | | | | |
| DAY 7 | | | | | |
| DAY 8 | | | | | |
| DAY 9 | | | | | 5. Pour off and discard all but 0.5 ml of the supernate. |
| DAY 10 | | | | | |
| DAY 11 | | | | | 6. Resuspend the sediment in the remaining 0.5 ml of supernate by touching the bottom of the tube to a vortex machine or by flicking the |
| DAY 12 | | | | | |
| DAY 13 DAY 14 | | | | | bottom of the tube with your finger. |
| DAY 14 DAY 15 | | | | | 7. Transfer a drop of the resuspended |
| DAT 15 DAY 16 | | | | | sediment to a clean dry microscope slide and cover with a cover slip. |
| DAT 10 DAY 17 | | | | | |
| DAT 17 DAY 18 | | | | | Count and record the <i>average</i> number of cells found in 10 high power fields. |
| DAY 19 | | | | | 9. At the end of the month, add the column of |
| DAY 20 | | | | | entries for MICROSCOPIC and/or SPECIFIC GRAVITY and enter the TOTAL at the bottom of the column. Determine the MEAN by |
| DAY 21 | | | | | |
| DAY 22 | | | | | dividing the TOTAL by the number of days the test was run. |
| DAY 23 | | | | | |
| DAY 24 | | | | | 10. Store at 2° - 8°C. May be stored at room temperature once bottle is in use. |
| DAY 25 | | | | | |
| DAY 26 | | | | | *NOTE:The value range for Alta's Microscopic Control is based on the parameters set forth in the |
| DAY 27 | | | | | above procedure. Laboratories using a procedure with different parameters (i.e. volume, rpm and time |
| DAY 28 | | | | | of centrifugation and amount of supernate discarded) should develop their own range of values |
| DAY 29 | | | | | and mean for the control using their procedure. |
| DAY 30 | | | | | |
| DAY 31 | | | | | |
| TOTAL | | | | | |
| MEAN | | | | | |